



**NATIONWIDE
CHILDREN'S**

When your child needs a hospital, everything matters.

Sudden Neurologic Deterioration (SND) & Suspected Stroke Inpatient

**Center for
Clinical Excellence**

Inclusion Criteria

- Patients with new-onset (<24 hrs) neurologic signs or symptoms concerning for stroke

Exclusion Criteria

- Patient in NICU

**New-onset
neurologic signs or symptoms
concerning for stroke
in hospitalized patient**

Bedside nurse pages **primary team
resident/APP to bedside** for
Focused History & Physical Exam

- Review **Risk Factors for Stroke**
- Discuss with primary Attending (or Safety Officer (SOD) if attending not available)

**Concern for stroke when
neurological signs or symptoms
are:**

- New (within 24 hours), sudden onset and focal

Signs or Symptoms may include:

- Unilateral weakness
- Speech difficulty (aphasia or dysarthria)
- Vision loss or double vision
- Ataxia
- Unilateral sensory change
- Vertigo
- New onset focal seizure with atypical prolonged postictal paralysis

Off Pathway

Review **Differential Diagnosis**

Primary team concerned for
potential stroke?

Yes

- Order **Neuroprotective care and labs.**
- Page on call Neurology Fellow** with "**urgent consult-stroke evaluation, Rm#, Call back #**" to meet primary team (senior resident) and bedside RN for **Bedside Huddle** within 30 min. Notify charge RN of stroke huddle.

Off Pathway

Continue diagnostic evaluation and
treatment as clinically indicated

Neurology concerned
for potential stroke?

Yes

SND – Inpatient Team Activation

- SND - Inpatient team activation per Neurology team member decision**
- Unit **charge RN calls** Emergency Communications Center (ECC) at 2-2233 to request "**SND Inpatient Team Activation**" (to start situational awareness for radiology and multidisciplinary stroke team)
- If patient has sickle cell disease, primary team to page hematology 1st on call for additional recommendations
- ACT as indicated

Stroke Evaluation

- Primary team orders **neuroimaging (MRI)** based on Neurology/ Inpatient SND Team recommendation
- CT head without contrast is NOT routinely recommended

Must NOTIFY radiology
(by activating **Inpatient
SND Alert** or by phone call)
in addition to placing STAT
MRI order.

Manage as indicated based on imaging
and clinical signs and symptoms

Imaging shows
likely stroke?

No

Yes

Individualized management
Off Pathway

- Individualized **stroke management** per Neurology and Neurosurgery recommendations in discussion with primary team and Inpatient SND Team
- ACT if patient on floor

Focused History & Physical Exam

HPI

- Any baseline developmental/cognitive delay/disability
- Any baseline neurological deficits
- Time of onset/last known well/baseline
- New-onset focal seizure
- Symptoms including:
 - Ataxia
 - Speech difficulty
 - Unilateral weakness
 - Vision change
 - Unilateral sensory change
 - Vertigo
- Symptoms improving or worsening
- Medications including recent chemotherapy
- Recent head injury or neck flexion/extension
- Recent illness especially head/neck infection
- Any [Risk Factors for stroke](#)

PMH

- Any [Risk Factors for stroke](#)
- Presence of hardware including dental

Family History

- Stroke or heart attack before the age of 50
- Leg or lung clots
- Multiple miscarriages
- Vasculopathy

Physical Exam

- [Peds NIH Stroke Scale, Use MDCalc](#)
- Other exam as clinically indicated

- If decreased alertness/responsiveness, consider [Glasgow Coma Scale, Use MDCalc](#)

[Return to Algorithm](#)

Risk Factors for Pediatric Stroke

- **Cardiac**
 - Congenital heart disease with a shunting lesion
 - Mechanical valve or mechanical assist device
 - Arrhythmias
 - Heart failure
 - Endocarditis
- **Hematology & Coagulation**
 - Sickle cell disease
 - Hypercoagulable condition including oral contraceptives with estrogen, recent treatment with L-Asparaginase
 - Dehydration
- **Cerebral vasculopathy & Vasculitis**
 - Moyamoya disease or systemic large vessel vasculitis, such as Takayasu's
- **Metabolic**
 - Mitochondrial encephalopathy
- **Genetic**
 - Trisomy 21
 - Neurofibromatosis
 - Connective tissue disorders
 - PHACE(S) Syndrome:
 - Posterior fossa malformations
 - Hemangioma of the cervicofacial region
 - Arterial anomalies
 - Cardiac anomalies
 - Eye anomalies
 - Sternal or abdominal clefting or ectopia cordis
- **Head, neck or CNS infections**
 - Meningitis/encephalitis
 - Mastoiditis
 - Lemierre's syndrome (septic thrombophlebitis of the internal jugular vein)
- **Trauma**
 - Recent minor head or neck trauma
 - Non-accidental trauma
- **Rheumatological disease**
 - Lupus
- **Medications**
 - Oral contraceptives with estrogen
 - L-Asparaginase
 - IVIG
- **Oncologic**

[Return to Algorithm](#)

Differential Diagnoses

Trauma	Vascular	Neoplastic
<ul style="list-style-type: none"> • Cerebral edema • Cerebral contusion • Intracranial hemorrhage • Extracranial hemorrhage 	<ul style="list-style-type: none"> • Strokes • Intracerebral hemorrhage • Dural sinus thrombosis • Vascular anomalies/malformation 	<ul style="list-style-type: none"> • Brain tumor • Chemotherapy toxicities
Inflammatory/Infection	Metabolic	Other
<ul style="list-style-type: none"> • Meningitis • Encephalitis • Abscess • Acute cerebellar ataxia • Cerebellitis • Demyelinating disease • Bell's Palsy 	<ul style="list-style-type: none"> • Inborn errors of metabolism • Hepatic encephalopathy • Renal failure • Hypoglycemia 	<ul style="list-style-type: none"> • Hydrocephalus • Idiopathic intracranial hypertension • Seizures/epilepsy • Migraine • Intoxication/drug toxicity • Conversion disorder

[Return to Algorithm](#)

“SND- Inpatient Team Activation” Process

Initial Bedside Huddle with:

- Primary team senior resident (and intern)
- Neurology fellow (team member)
- Bedside nurse
- Vascular Access Team (VAT) consult as indicated
- Optional: Unit Charge nurse

If patient has sickle cell disease, primary team will page hematology 1st on call

SND- Inpatient Team Activation

“SND- Inpatient Team Activation” Objectives

- To notify radiology of need for STAT brain MRI
- To notify multidisciplinary team of patient with potential need of STAT stroke care (ie. situational awareness of potential stroke in inpatient)

“SND- Inpatient Team Activation” Process

- SND- Inpatient Team Activation per **Neurology decision**
- Unit **charge nurse calls ECC** to request “*SND- Inpatient Team Activation*”
- ECC will page Inpatient SND Team members with “**SND- Inpatient Team Activation, pt age/gender, building/Unit/Rm#**”. Potential incoming orders. No response by team members required.

SND - Inpatient Team

For situational awareness of potential stroke in inpatient; no response required

- | | |
|---|--|
| • Safety Officer | • Nursing supervisor |
| • Neurology fellow | • PICU attending (H2B) |
| • Neurology attending (will not be paged) | • PICU fellow (H2B) |
| • Radiology / Neuroradiology | • PICU charge nurse |
| • MRI technician | • Neurosurgery |
| • Anesthesia | • Vascular Access Team (VAT) (not paged) |

[Return to Algorithm](#)

Diagnostic Testing & Neuroprotective Care

Diagnostic Testing

Labs:

- POC glucose (notify MD if <80 or >180)
- CBC with differential
- CMP
- PT/PTT/INR
- Fibrinogen
- Serum-HCG if indicated
- Type/screen
- If sickle cell diagnosis - Hgb electrophoresis

[Neuroimaging](#)

Optional labs:

- Urinalysis
- Urine drug screen
- Serum ethanol screening

Optional Imaging:

- CXR

Optional Other:

- EKG

Neuroprotective Care

- Vital signs Q30Min (resident to wean as indicated)
- Supplemental oxygen to maintain saturations $\geq 94\%$ (except if cyanotic heart disease)
- IV access x 2; large bore, prefer upper extremity
- Normal saline at maintenance rate
- NS bolus as clinically indicated
- NPO
- Bed rest
- Head of bed flat except if concern for increased intracranial pressure (ICP), then elevate head of bed 30 degrees
- Acetaminophen 15 mg/kg Rectal/IV/PO Q6H PRN temperature >37.5 C
- Notify resident of any seizure-like activity

[Return to Algorithm](#)

Neuroimaging

- **Primary team should confirm imaging with Neurology Fellow before ordering**
- **STAT MRI** (MR Head Short Stroke/Sudden Neurologic Deterioration)
 - If abnormal, reflex to MRA Head and MRA Neck, without Contrast, per Radiology
- If unable to obtain MRI (preferred modality):
 - CT Stroke-Sudden Neurologic Deterioration without contrast
 - CT Head Neck Angiography with contrast

- STAT MRI order will not be **automatically** scheduled without “**SND Inpatient Team Activation**” **through ECC**. Activation of the Inpatient SND Team by an ECC page will automatically schedule MRI, mobilize MRI team and notify neuroradiology. Neuroradiologist may reach out to the ordering provider or can be contacted at 2-9244 after SND Team initiated, but this is not required.
- Any stat MRI requests outside of the inpatient SND, especially after hours, must be discussed prior with the Radiologist by calling 2-9244

[Return to Algorithm](#)

Stroke Management

Patients with stroke require individualized management per Neurology recommendations.

- Ischemic stroke
 - Consider alteplase (tPA) within 4.5 hours
 - Thrombectomy for proximal large vessel occlusion
 - Antithrombotic therapy
 - Consider unfractionated heparin or LMW heparin (aka enoxaparin)
 - Consider aspirin
- Hemorrhagic stroke
 - Neurosurgery consult
- Stroke in sickle cell disease
 - Hematology consult
 - Consider simple vs. exchange transfusion

All other diagnoses should be treated using the standard of care in conjunction with appropriate consultation

[**Return to Algorithm**](#)

Deterioration & Escalation of Care

Identification of Deterioration

- Declining mental status
- Concerning vital signs (inc. Cushing triad: hypertension, bradycardia and irregular respiration, concerning for increased intracranial pressure)
- Worsening NIH Stroke Scale
- Status epilepticus
- Provider or parental concern

Escalation of Care

- ACT

[Return to Algorithm](#)

Severity Assessment

- [Pediatric NIH Stroke Scale](#)

- [Glasgow Coma Scale](#)

[Return to Algorithm](#)

Patient & Caregiver Education

- [Pediatric Stroke Family Tool Kit](#)
- **Helping Hands**
 - [Stroke in Children](#)
 - [Sickle Cell Disease and Stroke](#)

[Return to Algorithm](#)

Metrics

Pathway Goal

The Sudden Neurologic Deterioration & Suspected Stroke pathway will improve the timeliness of diagnosis in hospitalized patients with new-onset symptoms concerning for stroke.

Quality Measures

Outcome Metrics

- Primary Outcome metric:
 - Imaging within 1 hour of patient on pathway
- Specialty metrics:
 - Rate of missed stroke or other serious new-onset neurological condition for patient meeting pathway inclusion criteria, with or without activation of pathway
 - Rate of stroke or other new finding on diagnostic imaging for “stroke concern”

Process Metrics

- Order set use
- % of patients on pathway getting MRI vs. HCT
- Rate stroke of alerts activation

Balancing Metrics

- Rate of normal finding on diagnostic imaging for stroke concern

[Return to Algorithm](#)

References

1. Rivkin MJ, Bernard TJ, Dowling MM, Amlie-Lefond C. Corrigendum to 'Guidelines for Urgent Management of Stroke in Children' [Pediatric Neurology 56 (2016) 8-17]. *Pediatr Neurol*. 2016;64:105. doi:10.1016/j.pediatrneurol.2016.08.019
2. Powers WJ, Rabinstein AA, Ackerson T, et al. 2018 Guidelines for the Early Management of Patients With Acute Ischemic Stroke: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association [published correction appears in Stroke. 2018 Mar;49(3):e138. doi: 10.1161/STR.0000000000000163] [published correction appears in Stroke. 2018 Jun;49(6):e233-e234. doi: 10.1161/STR.0000000000000172]. *Stroke*. 2018;49(3):e46-e110. doi:10.1161/STR.0000000000000158
3. Ferriero DM, Fullerton HJ, Bernard TJ, et al. Management of Stroke in Neonates and Children: A Scientific Statement From the American Heart Association/American Stroke Association. *Stroke*. 2019;50(3):e51-e96. doi:10.1161/STR.0000000000000183
4. Teasdale G, Maas A, Lecky F, Manley G, Stocchetti N, Murray G. The Glasgow Coma Scale at 40 years: standing the test of time [published correction appears in Lancet Neurol. 2014 Sep;13(9):863]. *Lancet Neurol*. 2014;13(8):844-854. doi:10.1016/S1474-4422(14)70120-6
5. Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet*. 1974;2(7872):81-84. doi:10.1016/s0140-6736(74)91639-0
6. Beslow LA, Kasner SE, Smith SE, et al. Concurrent validity and reliability of retrospective scoring of the Pediatric National Institutes of Health Stroke Scale. *Stroke*. 2012;43(2):341-345. doi:10.1161/STROKEAHA.111.633305

[Return to Algorithm](#)

Team & Process

Pathway Development Team

Leader(s):

Hospital Pediatrics:

Mathew Suer, MD

Neurology:

Melissa Chung, MD

Members:

Neurology:

Shawn Aylward, MD

Alyssa-Claire Mendoza, MD

Neurosurgery:

Eric Sribnick, MD

Radiology:

Jeremy Jones, MD

Christopher Gerity, MPH, RT

PICU:

Jennifer MacDonald, MD

Marlina Lovett, MD

Resident:

Meaghan Holzworth, MD

Nursing:

Kathy Moellman, RN

Pharmacy:

Patrik Miller, PharmD

Clinical Pathways Program:

Medical Director – Hospital Pediatrics:

Gerd McGwire, MD, PhD

Medical Director – Clinical Informatics & Emergency Medicine:

Laura Rust, MD, MPH

Business & Development Manager:

Rekha Voruganti, MBOE, LSSBB

Program Coordinators:

Tara Dinh, BS

Clinical Pathway Approved

Medical Director – Associate Chief Quality Officer, Center for Clinical Excellence:

Ryan Bode, MD, MBOE

Origination Date: *August, 2024*

Next Revision Date: *January 2028*

Clinical Pathway Development

This clinical pathway was developed using the process described in the NCH Clinical Pathway Development Manual Version 6, 2022. Clinical Pathways at Nationwide Children's Hospital (NCH) are standards which provide general guidance to clinicians. Patient choice, clinician judgment, and other relevant factors in diagnosing and treating patients remain central to the selection of diagnostic tests and therapy. The ordering provider assumes all risks associated with care decisions. NCH assumes no responsibility for any adverse consequences, errors, or omissions that may arise from the use or reliance on these guidelines. NCH's clinical pathways are reviewed periodically for consistency with new evidence; however, new developments may not be represented, and NCH makes no guarantees, representations, or warranties with respect to the information provided in this clinical pathway.

Copyright © 2023. Nationwide Children's Hospital. All rights reserved. No part of this document may be reproduced, displayed, modified, or distributed in any form without the express written permission of Nationwide Children's Hospital.

**For more information about our pathways and program please contact:
ClinicalPathwaysProgram@NationwideChildrens.org**

[Return to Algorithm](#)

NIH Stroke Scale

National Institutes of Health Stroke Scale score

1a. Level of consciousness	0 = Alert; keenly responsive 1 = Not alert, but arousable by minor stimulation 2 = Not alert; requires repeated stimulation 3 = Unresponsive or responds only with reflex
1b. Level of consciousness questions: What is the month? What is your age?	0 = Answers two questions correctly 1 = Answers one question correctly 2 = Answers neither question correctly
1c. Level of consciousness commands: Open and close your eyes. Grip and release your hand.	0 = Performs both tasks correctly 1 = Performs one task correctly 2 = Performs neither task correctly
2. Best gaze	0 = Normal 1 = Partial gaze palsy 2 = Forced deviation
3. Visual	0 = No visual loss 1 = Partial hemianopia 2 = Complete hemianopia 3 = Bilateral hemianopia
4. Facial palsy	0 = Normal symmetric movements 1 = Minor paralysis 2 = Partial paralysis 3 = Complete paralysis of one or both sides
5. Motor arm 5a. Left arm 5b. Right arm	0 = No drift 1 = Drift 2 = Some effort against gravity 3 = No effort against gravity; limb falls 4 = No movement
6. Motor leg 6a. Left leg 6b. Right leg	0 = No drift 1 = Drift 2 = Some effort against gravity 3 = No effort against gravity 4 = No movement
7. Limb ataxia	0 = Absent 1 = Present in one limb 2 = Present in two limbs
8. Sensory	0 = Normal; no sensory loss 1 = Mild-to-moderate sensory loss 2 = Severe to total sensory loss
9. Best language	0 = No aphasia; normal 1 = Mild to moderate aphasia 2 = Severe aphasia 3 = Mute, global aphasia
10. Dysarthria	0 = Normal 1 = Mild to moderate dysarthria 2 = Severe dysarthria
11. Extinction and inattention	0 = No abnormality 1 = Visual, tactile, auditory, spatial, or personal inattention 2 = Profound hemi-inattention or extinction

Total score = 0–42.

[Peds NIH Stroke Scale with instructions](#)

[Return to Algorithm](#)

Pediatric NIH Stroke Scale with Instructions

Item# and Instructions	Scale Definition and Scoring Guide
1a. Level of Consciousness: the investigator must choose a response, even if a full evaluation is prevented by such obstacles as an endotracheal tube, language barrier, orotracheal trauma/bandages. A 3 is scored only if the patient makes no movement (other than reflexive posturing) in response to noxious stimulation.	0 = Alert; keenly responsive. 1 = Not alert, but arousable by minor stimulation to obey, answer, or respond. 2 = Not alert, requires repeated stimulation to attend, or is obtunded and requires strong or painful stimulation to make movements (not stereotyped). 3 = Responds only with reflex motor or autonomic effects or totally unresponsive, flaccid, areflexic.
b. LOC Questions: The patient is asked the month and his/her age. The answer must be correct - there is no partial credit for being close. Aphasic and stuporous patients who do not comprehend the questions will score 2. Patients unable to speak because of endotracheal intubation, orotracheal trauma, severe dysarthria from any cause, language barrier or any other problem not secondary to aphasia are given a 1. It is important that only the initial answer be graded and that the examiner not "help" the patient with verbal or non-verbal cues. <i>Modified for children, age 2 years and up. A familiar Family Member must be present for this item: Ask the child "how old are you?" Or "How many years old are you?" for question number one. Give credit if the child states the correct age, or shows the correct number of fingers for his/her age. For the second question, ask the child "where is XX?", XX referring to the name of the parent or other familiar family member present. Use the name for that person which the child typically uses, e.g. "mommy". Give credit if the child correctly points to or gazes purposefully in the direction of the family member.</i>	0 = Answers both questions correctly. 1 = Answers one question correctly. 2 = Answers neither question correctly.
1c. LOC Commands: The patient is asked to open and close the eyes and then to grip and	0 = Performs both tasks correctly 1 = Performs one task correctly

[Return to Algorithm](#)

release the non-paretic hand. <i>For children one may substitute the command to grip the hand with the command "show me your nose" or "touch your nose".</i> Substitute another one step command if the hands cannot be used. Credit is given if an unequivocal attempt is made but not completed due to weakness. If the patient does not respond to command, the task should be demonstrated to them (pantomime) and score the result (i.e., follows none, one or two commands). Patients with trauma, amputation, or other physical impediments should be given suitable one-step commands. Only the first attempt is scored.	2 = Performs neither task correctly
2. Best Gaze: Only horizontal eye movements will be tested. Voluntary or reflexive (oculocephalic) eye movements will be scored but caloric testing is not done. If the patient has a conjugate deviation of the eyes that can be overcome by voluntary or reflexive activity, the score will be 1. If a patient has an isolated peripheral nerve paresis (CN III, IV or VI) score a 1. Gaze is testable in all aphasic patients. Patients with ocular trauma, bandages, preexisting blindness or other disorder of visual acuity or fields should be tested with reflexive movements and a choice made by the investigator. Establishing eye contact and then moving about the patient from side to side will occasionally clarify the presence of a partial gaze palsy.	0 = Normal 1 = Partial gaze palsy. This score is given when gaze is abnormal in one or both eyes, but where forced deviation or total gaze paresis are not present. 2 = Forced deviation, or total gaze paresis not overcome by the oculocephalic maneuver.
3. Visual: Visual fields (upper and lower quadrants) are tested by confrontation, using finger counting (<i>for children > 6 years</i>) or visual threat (<i>for children 2 to 6 years</i>) as appropriate. Patient must be encouraged, but if they look at the side of the moving fingers appropriately, this can be scored as normal. If there is unilateral blindness or enucleation, visual fields in the remaining eye are scored. Score 1 only if a clear-cut asymmetry, including quadrantanopia is found. If patient is blind from any cause score 3. Double simultaneous stimulation is performed at this	0 = No visual loss 1 = Partial hemianopia 2 = Complete hemianopia 3 = Bilateral hemianopia (blind including cortical blindness)

[Return to Algorithm](#)

point. If there is extinction patient receives a 1 and the results are used to answer question 11.	
4. Facial Palsy: Ask, or use pantomime to encourage the patient to show teeth or raise eyebrows and close eyes. Score symmetry of grimace in response to noxious stimuli in the poorly responsive or non-comprehending patient. If facial trauma/bandages, orotracheal tube, tape or other physical barrier obscures the face, these should be removed to the extent possible.	0 = Normal symmetrical movement 1 = Minor paralysis (flattened nasolabial fold, asymmetry on smiling) 2 = Partial paralysis (total or near total paralysis of lower face) 3 = Complete paralysis of one or both sides (absence of facial movement in the upper and lower face)
5 & 6. Motor Arm and Leg: The limb is placed in the appropriate position: extend the arms (palms down) 90 degrees (if sitting) or 45 degrees (if supine) and the leg 30 degrees (always tested supine). Drift is scored if the arm falls before 10 seconds or the leg before 5 seconds. <i>For children too immature to follow precise directions or uncooperative for any reason, power in each limb should be graded by observation of spontaneous or elicited movement according to the same grading scheme, excluding the time limits.</i> The aphasic patient is encouraged using urgency in the voice and pantomime but not noxious stimulation. Each limb is tested in turn, beginning with the nonparetic arm. Only in the case of amputation or joint fusion at the shoulder or hip, or immobilization by an IV board, may the score be "9" and the examiner must clearly write the explanation for scoring as a "9". Score each limb separately.	5a. Left Arm 5b. Right Arm 0 = No drift, limb holds 90 (or 45) degrees for full 10 seconds. 1 = Drift, Limb holds 90 (or 45) degrees, but drifts down before full 10 seconds; does not hit bed or other support. 2 = Some effort against gravity, limb cannot get to or maintain (if cued) 90 (or 45) degrees, drifts down to bed, but has some effort against gravity. 3 = No effort against gravity, limb falls. 4 = No movement 9 = Amputation, joint fusion explain: 6a. Left Leg 6b. Right Leg 0 = No drift, leg holds 30 degrees position for full 5 seconds. 1 = Drift, leg falls by the end of the 5 second period but does not hit bed. 2 = Some effort against gravity; leg falls to bed by 5 seconds, but has some effort against gravity. 3 = No effort against gravity, leg falls to bed immediately. 4 = No movement 9 = Amputation, joint fusion explain:
7. Limb Ataxia: This item is aimed at finding evidence of a unilateral cerebellar lesion. Test with eyes open. In case of visual defect, insure testing is done in intact visual field. The fingernose-finger and heel-shin tests are performed on both sides, and ataxia is scored only if present out of proportion to weakness. <i>In children, substitute this task with reaching for a toy for the upper extremity, and kicking a toy or the examiner's hand, in children too</i>	0 = Absent 1 = Present in one limb 2 = Present in two limbs

[Return to Algorithm](#)

<i>young (< 5 years) or otherwise uncooperative for the standard exam item.</i> Ataxia is absent in the patient who cannot understand or is paralyzed. Only in the case of amputation or joint fusion may the item be scored "9", and the examiner must clearly write the explanation for not scoring. In case of blindness test by touching nose from extended arm position.	
8. Sensory: Sensation or grimace to pin prick when tested, or withdrawal from noxious stimuli in the obtunded or aphasic patient. <i>For children too young or otherwise uncooperative for reporting gradations of sensory loss, observe for any behavioral response to pin prick, and score it according to the same scoring scheme as a "normal" response, "mildly diminished" or "severely diminished" response.</i> Only sensory loss attributed to stroke is scored as abnormal and the examiner should test as many body areas [arms (not hands), legs, trunk, face] as needed to accurately check for hemisensory loss. A score of 2, "severe or total," should only be given when a severe or total loss of sensation can be clearly demonstrated. Stuporous and aphasic patients will therefore probably score 1 or 0.	0 = Normal; no sensory loss. 1 = Mild to moderate sensory loss; patient feels pinprick is less sharp or dull on the affected side; or there is a loss of superficial pain with pinprick but patient is aware he/she is being touched. 2 = Severe to total sensory loss; patient is not aware of being touched in the face, arm, and leg.
9. Best Language: A great deal of information about comprehension will be obtained during the preceding sections of the examination. <i>For children age 6 years and up with normal language development before onset of stroke: The patient is asked to describe what is happening in the attached picture, to name the items on the attached naming sheet, to repeat words from the attached list of sentences (Table S1; Fig S1, S2, S3).</i> Comprehension is judged from responses here as well as to all of the commands in the preceding general neurological exam. If visual loss interferes with the tests, ask the patient to identify objects placed in the hand, repeat, and produce speech. The intubated patient should be asked to write. The patient in coma (question 1a=3) will arbitrarily score	0 = No aphasia, normal 1 = Mild to moderate aphasia; some obvious loss of fluency or facility of comprehension, without significant limitation on ideas expressed or form of expression. Reduction of speech and/or comprehension, however, makes conversation about provided materials difficult or impossible. For example in conversation about provided materials examiner can identify picture or naming card from patient's response. 2 = Severe aphasia; all communication is through fragmentary expression; great need for inference, questioning, and guessing by the listener. Range of information that can be exchanged is limited; listener carries burden of communication. Examiner cannot identify materials provided from patient response.

[Return to Algorithm](#)

3 on this item. The examiner must choose a score in the patient with stupor or limited cooperation but a score of 3 should be used only if the patient is mute and follows no one step commands. <i>For children age 2 yrs to 6 yrs (or older children with premorbid language skills < 6 yr level), score this item based on observations of language comprehension and speech during the examination.</i>	3 = Mute, global aphasia; no usable speech or auditory comprehension.
10. Dysarthria: If patient is thought to be normal an adequate sample of speech must be obtained by adequate patient to read or repeat words from the attached list. If the patient has severe aphasia, the clarity of articulation of spontaneous speech can be rated. Only if the patient is intubated or has other physical barrier to producing speech, may the item be scored "9", and the examiner must clearly write an explanation for not scoring. Do not tell the patient why he/she is being tested.	0 = Normal 1 = Mild to moderate; patient slurs at least some words and, at worst, can be understood with some difficulty. 2 = Severe; patient's speech is so slurred as to be unintelligible in the absence of or out of proportion to any dysphasia, or is mute/anarthric. 9 = Intubated or other physical barrier, explain:
11. Extinction and Inattention (formerly Neglect): Sufficient information to identify neglect may be obtained during the prior testing. If the patient has a severe visual loss preventing visual double simultaneous stimulation, and the cutaneous stimuli are normal, the score is normal. If the patient has aphasia but does appear to attend to both sides, the score is normal. The presence of visual spatial neglect or anosagnosia may also be taken as evidence of abnormality. Since the abnormality is scored only if present, the item is never untestable.	0 = No abnormality. 1 = Visual, tactile, auditory, spatial, or personal inattention or extinction to bilateral simultaneous stimulation in one of the sensory modalities. 2 = Profound hemi-inattention or hemi-inattention to more than one modality. Does not recognize own hand or orients to only one side of space.

[Return to Algorithm](#)

Glasgow Coma Scale

[The Glasgow Structured Approach to Assessment of the Glasgow Coma Scale](#)

Behavior	Response	Score
Eyes	Spontaneously	4
	To speech	3
	To Pain	2
	No response	1
Verbal	Oriented to time, place and person	5
	Confused	4
	Inappropriate words	3
	Incomprehensible sounds	2
	No response	1
Motor	Obeys commands	6
	Moves to localized pain	5
	Flexion with drawal from pain	4
	Abnormal flexion	3
	Abnormal extension	2
	No response	1
Total Score	Best Response	15
	Totally unresponsive	3

Teasdale et al. 2014
Teasdale et al. 1974

[Return to Algorithm](#)