

EXTERNAL WEBPAGE MASTER FORMULA

Compound Name: Quinidine	Container-closure system(s): tight, light resistant plastic container (Amber plastic bottle)
Strength: 10 mg/mL	Preservatives: Contained in suspending agent or sweetener
Dosage Form: Oral Suspension	Beyond Use Date (compound type): 35 days
Product Description: opaque suspension	Storage: Refrigerate (Preferred) or Room Temperature
Auxiliary Labels: <input type="checkbox"/> Take with food <input checked="" type="checkbox"/> Shake Well <input type="checkbox"/> For external use only <input type="checkbox"/> Refrigerate <input type="checkbox"/> Do not use after: <input type="checkbox"/> Do not take with dairy products <input type="checkbox"/> Do not refrigerate <input type="checkbox"/> For the Ears <input type="checkbox"/> Avoid prolonged sun exposure <input type="checkbox"/> Take even if you feel fine <input type="checkbox"/> For the Eyes <input checked="" type="checkbox"/> Other: Protect from light	
Quality control procedures (ex: pH test, etc.): Visual inspection of final product	
Ingredients: Quinidine Sulfate 200 mg tablet Ora-Blend (QS)	

Instructions for preparation:**Calculations:****Directions:**

1. Crush tablets into a fine powder using a mortar & pestle.
2. Wet powder with minimal amount of Ora-Blend. Levigate to a uniform paste.
3. Add vehicle in geometric proportions almost to final volume.
4. Pour mixture into graduated cylinder.
5. Rinse mortar with remaining vehicle. Add to graduated cylinder.
6. QS to final volume. Mix well.

Ingredient Name/Strength/Form	QS	Quantity	Units
Quinidine Sulfate 200 mg tablets		6	Tablets
Ora Blend	X	120	mL

Equipment/ Supplies:

- | | | |
|---|--|---|
| <input checked="" type="checkbox"/> Mortar and Pestle | <input checked="" type="checkbox"/> Stirring rod | <input type="checkbox"/> Ointment slab |
| <input checked="" type="checkbox"/> Graduated cylinder(s) | <input type="checkbox"/> Spatula | <input checked="" type="checkbox"/> Balance |
| <input type="checkbox"/> Weigh paper | <input checked="" type="checkbox"/> Amber (plastic/glass) bottle | <input type="checkbox"/> Ointment jar |

Revised: TS 11/25



NATIONWIDE CHILDREN'S®
When your child needs a hospital, everything matters.

☒ Counting tray

☐ Syringe (Oral)

☐ Funnel

References:

1. Allen LV Jr, Erickson MA III. Stability of bethanechol chloride, pyrazinamide, quinidine sulfate, rifampin, and tetracycline hydrochloride in extemporaneously compounded oral liquids. *Am J Health- Syst Pharm.* 1998; 55:1804-1809
2. US Pharmacopeial Convention, Inc. United States Pharmacopeia 42. Chapter <795> “Pharmaceutical Compounding–Non- Sterile Preparations.” Rockville, MD: US Pharmacopeial Convention, Inc.; 2022

Nationwide Children’s Hospital Disclaimer:

The information provided on this site is for educational and informational purposes and may not be relied upon to direct or provide care to a patient. This material is not intended as a substitute for consulting health care professionals. The health care professionals remain solely responsible to provide and direct patient care. The information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professional or pharmacist. Nationwide Children's Hospital assumes no liability for any injury and/or damage to persons from use of any information contained on this site.

Revised: TS 11/25

